

Contents

Volume 107 1989

Foe, V. E. Mitotic domains reveal early commitment of cells in <i>Drosophila</i> embryos	1
Stirpe, N. S. and Goetinck, P. F. Gene regulation during cartilage differentiation: temporal and spatial expression of link protein and cartilage matrix protein in the developing limb	23
Callaini, G. Microtubule distribution reveals superficial metameric patterns in the early <i>Drosophila</i> embryo	35
Messenger, N. J. and Warner, A. E. The appearance of neural and glial cell markers during early development of the nervous system in the amphibian embryo	43
Tam, P. P. L. Regionalisation of the mouse embryonic ectoderm: allocation of prospective ectodermal tissues during gastrulation	55
Saló, E. and Baguñà, J. Regeneration and pattern formation in planarians. II. Local origin and role of cell movements in blastema formation	69
Baguñà, J., Saló, E. and Auladell, C. Regeneration and pattern formation in planarians. III. Evidence that neoblasts are totipotent stem cells and the source of blastema cells	77
Crandall, I. E. and Newell, P. C. Changes in cell surface glycoproteins during <i>Dictyostelium</i> development analysed using monoclonal antibodies	87
Taketo-Hosotani, T., Nishioka, Y., Nagamine, C. M., Villalpando, I. and Merchant-Larios, H. Development and fertility of ovaries in the B6.Y ^{DOM} sex-reversed female mouse	95
Eccleston, P. A., Bannerman, P. G. C., Pleasure, D. E., Winter, J., Mirsky, R. and Jessen, K. R. Control of peripheral glial cell proliferation: enteric neurons exert an inhibitory influence on Schwann cell and enteric glial cell DNA synthesis in culture	107
Heath, J. K., Paterno, G. D., Lindon, A. C. and Edwards, D. R. Expression of multiple heparin-binding growth factor species by murine embryonal carcinoma and embryonic stem cells	113
Cuthbertson, R. A., Beck, F., Senior, P. V., Haralambidis, J., Penschow, J. D. and Coghlan, J. P. Insulin-like growth factor II may play a local role in the regulation of ocular size	123
Gaunt, S. J., Krumlauf, R. and Duboule, D. Mouse homeo-genes within a subfamily, Hox-1.4, -2.6 and -5.1, display similar anteroposterior domains of expression in the embryo, but show stage- and tissue-dependent differences in their regulation	131
Carr, J. N. and Taghert, P. H. Pair-rule expression of a cell surface molecule during gastrulation of the moth embryo	143
Srinivas, U. K. and Henderson, E. J. Biochemical differentiation in a mutant of <i>Dictyostelium discoideum</i> defective in cyclic AMP chemotaxis and in intercellular cohesion	153
Sapienza, C., Paquette, J., Tran, T. H. and Peterson, A. Epigenetic and genetic factors affect transgene methylation imprinting	165
ESSAY IN DEVELOPMENT	
Del Pino, E. M. Modifications of oogenesis and development in marsupial frogs	169
CONTRIBUTED PAPERS	
Grossniklaus, U., Bellen, H. J., Wilson, C. and Gehring, W. J. P-element-mediated enhancer detection applied to the study of oogenesis in <i>Drosophila</i>	189

Contents

Patel, N. H., Kornberg, T. B. and Goodman, C. S. Expression of <i>engrailed</i> during segmentation in grasshopper and crayfish	201	Oppenheim, R. W., Bursztajn, S. and Prevet, D. Cell death of motoneurons in the chick embryo spinal cord. XI. Acetylcholine receptors and synaptogenesis in skeletal muscle following the reduction of motoneuron death by neuromuscular blockade	331
Kuopio, T., Tapanainen, J., Pelliniemi, L. J. and Huhtaniemi, I. Developmental stages of fetal-type Leydig cells in prepubertal rats	213	Galliot, B., Dollé, P., Vigneron, M., Featherstone, M. S., Baron, A. and Duboule, D. The mouse Hox-1.4 gene: primary structure, evidence for promoter-activity and expression during development	343
McAvoy, J. W. and Chamberlain, C. G. Fibroblast growth factor (FGF) induces different responses in lens epithelial cells depending on its concentration	221	Rudnicki, M. A., Reuhl, K. R. and McBurney, M. W. Cell lines with developmental potential restricted to mesodermal lineages isolated from differentiating cultures of pluripotent P19 embryonal carcinoma cells	361
Cooke, J. Mesoderm-inducing factors and Spemann's organiser phenomenon in amphibian development	229	Sutcliffe, M. J. and Burgoyne, P. S. Analysis of the testes of H-Y negative XOSr ^b mice suggests that the spermatogenesis gene (<i>Spy</i>) acts during the differentiation of the A spermatogonia	373
Duxson, M. J. and Usson, Y. Cellular insertion of primary and secondary myotubes in embryonic rat muscles	243	Cohen, J., Nurcombe, V., Jeffrey, P. and Edgar, D. Developmental loss of functional laminin receptors on retinal ganglion cells is regulated by their target tissue, the optic tectum	381
Fellah, J. S., Vault, D., Tournefier, A. and Charlemagne, J. Ontogeny of immunoglobulin expression in the Mexican axolotl	253	Hartenstein, V. and Posakony, J. W. Development of adult sensilla on the wing and notum of <i>Drosophila melanogaster</i>	389
Vandenbunder, B., Pardanaud, L., Jaffredo, T., Mirabel, M. A. and Stehelin, D. Complementary patterns of expression of <i>c-ets 1</i> , <i>c-myb</i> and <i>c-myc</i> in the blood-forming system of the chick embryo	265	Tsunoda, Y., Tokunaga, T., Imai, H. and Uchida, T. Nuclear transplantation of male primordial germ cells in the mouse	407
Mitrani, E. and Shimoni, Y. Retinoic acid inhibits growth in agarose of early chick embryonic cells and may be involved in regulation of axis formation	275	Fontaine-Péru, J. C., Chanconie, M., Le Douarin, N. M., Gershon, M. D. and Rothman, T. P. Mitogenic effect of muscle on the neuroepithelium of the developing spinal cord	413
Halfter, W. Antisera to basal lamina and glial endfeet disturb the normal extension of axons on retina and pigment epithelium basal laminae	281	Mitani, S. Retarded gastrulation and altered subsequent development of neural tissues in heparin-injected <i>Xenopus</i> embryos	423
Stephens, L., Kitajima, T. and Wilt, F. Autonomous expression of tissue-specific genes in dissociated sea urchin embryos	299	Birk, D. E., Southern, J. F., Zycband, E. I., Fallon, J. T. and Trelstad, R. L. Collagen fibril bundles: a branching assembly unit in tendon morphogenesis	437
Stern, C. D., Norris, W. E., Bronner-Fraser, M., Carlson, G. J., Faissner, R. J. and Schachner, M. J1/tenascin-related molecules are not responsible for the segmented pattern of neural crest cells or motor axons in the chick embryo	309		
Sánchez-Herrero, E. and Akam, M. Spatially ordered transcription of regulatory DNA in the bithorax complex of <i>Drosophila</i>	321		

Ohinata, H., Tochinal, S. and Katagiri, C. Ontogeny and tissue distribution of leukocyte-common antigen bearing cells during early development of <i>Xenopus laevis</i>	445	Hardy, K., Handyside, A. H. and Winston, R. M. L. The human blastocyst: cell number, death and allocation during late preimplantation development <i>in vitro</i>	597
Steel, K. P. and Barkway, C. Another role for melanocytes: their importance for normal stria vascularis development in the mammalian inner ear	453	Inouye, K. Control of cell type proportions by a secreted factor in <i>Dictyostelium discoideum</i>	605
West, J. D. and Flockhart, J. H. Genetic differences in glucose phosphate isomerase activity among mouse embryos	465	De, S. K., McMaster, M. T., Dey, S. K. and Andrews, G. K. Cell-specific metallothionein gene expression in mouse decidua and placenta	611
Moskalewski, S. and Malejczyk, J. Bone formation following intrarenal transplantation of isolated murine chondrocytes: chondrocyte – bone cell transdifferentiation	473	Kopczynski, C. C. and Muskavitch, M. A. T. Complex spatio-temporal accumulation of alternative transcripts from the neurogenic gene <i>Delta</i> during <i>Drosophila</i> embryogenesis	623
McConnell, J. and Lee, M. Presence of <i>cdc2</i> ⁺ -like proteins in the preimplantation mouse embryo	481	Brown, N. A., Hoyle, C. I., McCarthy, A. and Wolpert, L. The development of asymmetry: the sidedness of drug-induced limb abnormalities is reversed in situs inversus mice	637
Hooper, K. L., Parkhurst, S. M. and Ish-Horowicz, D. Spatial control of <i>hairy</i> protein expression during embryogenesis	489	McMahon, J. A. and McMahon, A. P. Nucleotide sequence, chromosomal localization and developmental expression of the mouse <i>int-1</i> -related gene	643
Nothiger, R., Jonglez, M., Leuthold, M., Meier-Gerschwiler, P. and Weber, T. Sex determination in the germ line of <i>Drosophila</i> depends on genetic signals and inductive somatic factors	505	Gaul, U. and Jäckle, H. Analysis of maternal effect mutant combinations elucidates regulation and function of the overlap of <i>hunchback</i> and <i>Krüppel</i> gene expression in the <i>Drosophila</i> blastoderm embryo	651
Storey, K. G. Cell lineage and pattern formation in the earthworm embryo	519	Vavra, S. H. and Carroll, S. B. The zygotic control of <i>Drosophila</i> pair-rule gene expression. I. A search for new pair-rule regulatory loci	663
Storey, K. G. The effects of ectoteloblast ablation in the earthworm embryo	533	Carroll, S. B. and Vavra, S. H. The zygotic control of <i>Drosophila</i> pair-rule gene expression. II. Spatial repression by gap and pair-rule gene products	673
Govind, C. K. and Pearce, J. Delayed determination of claw laterality in lobsters following loss of target	547	ESSAY IN DEVELOPMENT	
Easter, S. S. and Taylor, J. S. H. The development of the <i>Xenopus</i> retinofugal pathway: optic fibers join a pre-existing tract	553	Smith, L. D. The induction of oocyte maturation: transmembrane signaling events and regulation of the cell cycle	685
Busturia, A., Casanova, J., Sánchez-Herrero, E., González, R. and Morata, G. Genetic structure of the <i>abd-A</i> gene of <i>Drosophila</i>	575	CONTRIBUTED PAPERS	
Schaart, G., Viebahn, C., Langmann, W. and Ramaekers, F. Desmin and titin expression in early postimplantation mouse embryos	585	Sharpe, C. R., Pluck, A. and Gurdon, J. B. XIF3, a <i>Xenopus</i> peripherin gene, requires an inductive signal for enhanced expression in anterior neural tissue	701

Contents

Simcox, A. A., Roberts, I. J. H., Hersperger, E., Gribbin, M. C., Shearn, A. and Whittle, J. R. S. Imaginal discs can be removed from cultured embryos mutant for the segment-polarity genes <i>engrailed</i> , <i>naked</i> and <i>patched</i> but not from <i>wingless</i>	715	Mohun, T. J., Garrett, N. and Taylor, M. V. Temporal and tissue-specific expression of the proto-oncogene <i>c-fos</i> during development in <i>Xenopus laevis</i>	835
Basler, K. and Hafen, E. Dynamics of <i>Drosophila</i> eye development and temporal requirements of <i>sevenless</i> expression	723	Lawrence, P. A. and Johnston, P. Analysis of function of the pair-rule genes <i>hairy</i> , <i>even-skipped</i> and <i>fushi tarazu</i> in mosaic <i>Drosophila</i> embryos	847
Kobayashi, S. and Okada, M. Restoration of pole-cell-forming ability to u.v.-irradiated <i>Drosophila</i> embryos by injection of mitochondrial lrRNA	733	Tix, S., Bate, M. and Technau, G. M. Pre-existing neuronal pathways in the leg imaginal discs of <i>Drosophila</i>	855
Duxson, M. J., Usson, Y. and Harris, A. J. The origin of secondary myotubes in mammalian skeletal muscles: ultrastructural studies	743	Eichele, G. Retinoic acid induces a pattern of digits in anterior half wing buds that lack the zone of polarizing activity	863
Harris, A. J., Fitzsimons, R. B. and McEwan, J. C. Neural control of the sequence of expression of myosin heavy chain isoforms in foetal mammalian muscles	751	Lorenz, L. J., Hall, J. C. and Rosbash, M. Expression of a <i>Drosophila</i> mRNA is under circadian clock control during pupation	869
Harris, A. J., Duxson, M. J., Fitzsimons, R. B. and Rieger, F. Myonuclear birthdates distinguish the origins of primary and secondary myotubes in embryonic mammalian skeletal muscles	771	Boocock, C. A. Unidirectional displacement of cells in fibrillar matrices	881
Jones, E. A. and Woodland, H. R. Spatial aspects of neural induction in <i>Xenopus laevis</i>	785	Wilcox, M., DiAntonio, A. and Leptin, M. The function of PS integrins in <i>Drosophila</i> wing morphogenesis	891
Van Straaten, H. W. M., Hekking, J. W. M., Beursgens, J. P. W. M., Terwindt-Rouwenhorst, E. and Drukker, J. Effect of the notochord on proliferation and differentiation in the neural tube of the chick embryo	793	Kloc, M., Miller, M., Carrasco, A. E., Eastman, E. and Etkin, L. The maternal store of the <i>xlgr7</i> mRNA in full-grown oocytes is not required for normal development in <i>Xenopus</i>	899
Richardson, M. K., Hornbruch, A. and Wolpert, L. Pigment pattern expression in the plumage of the quail embryo and the quail-chick chimaera	805	Levi, G. and Teichberg, V. I. Patterns of expression of a 15K β -D-galactosidase-specific lectin during early development of the avian embryo	909
Barton, P. J. R., Harris, A. J. and Buckingham, M. E. Myosin light chain gene expression in developing and denervated fetal muscle in the mouse	819	Kam, E. and Pitts, J. D. Tissue-specific regulation of junctional communication in the skin of mouse fetuses homozygous for the repeated epilation (<i>Er</i>) mutation	923
Cameron-Curry, P., Dulac, C. and Le Douarin, N. M. Expression of the SMP antigen by oligodendrocytes in the developing avian central nervous system	825	Bagnall, K. M., Higgins, S. J. and Sanders, E. J. The contribution made by cells from a single somite to tissues within a body segment and assessment of their integration with similar cells from adjacent segments	931

**Dooley, T. P., Miranda, M., Jones, N. C.
and DePamphilis, M. L.**

Transactivation of the adenovirus EIIa promoter
in the absence of adenovirus E1a protein is
restricted to mouse oocytes and preimplantation
embryos

945

Index of Authors and Titles

957

Subject Index

961